

## DEGRADATION STUDY OF THIOPHANATE-METHYL RESIDUES IN CUCUMBER (*Cucumis sativus*)

<sup>1</sup>Ali Bassam Al-Obaidie, <sup>2</sup>Abdulkareem Jawad Ali, <sup>3</sup>Saleh Hassan Sumir,  
<sup>4</sup>Osama Imad Al-Samaraie, <sup>5</sup>Methal Hadi Ali

<sup>1,3</sup>Ministry of Higher Education and Scientific Research, Baghdad, Iraq. <sup>2,4,5</sup>National Center for Pesticides Control, Ministry of Agriculture, Baghdad, Iraq. E. mail: omertofe@yahoo.com

Article received 20.7.2018, Revised 15.9.2018, Accepted 22.9.2018

### ABSTRACT


We carried out the experiment under greenhouse conditions in plant protection department/ Ministry of Agriculture during 2017–2018 season, to assess the degradation sample of Thiophanate methyl (70% WP) fungicide in cucumber at three use rates (1.5gm/L, 3gm/L, and double spray 1.5gm/L). The researchers carried out the quantitative analysis of the fungicide residues as carbendazim was using High-performance liquid chromatography (HPLC) using QuEChERS extraction method. I found the rate of recovery was 92-106% and the comparative standard variation (RSD) were below 3.8%. Next the first order kinetics the fungicide degrades in cucumber i found the half life value in cucumber between 12.2–13.4 days unheeding of position and dose.

Keywords: Thiophanate-methyl residue - Degradation - Cucumber- QuEChERS – HPLC.

### INTRODUCTION

Toxic substances released from fungicides industries are One source of environmental pollution (Naqvi, et al., 2006) Thiophanate methyl [1, 2-bis-(3-methoxy carboxy-2-thioureidoben-zene)] is a large spectrum systemic fungicide which is extreme applied to control the major fungal diseases of different harvest [Maranghi, et al., 2003]. This-fungicides has ability to absorbed quickly by the plant during spraying it upon them. The compound can easily transport through parts of the plant as methyl benzimidazole carbamates (MBC) which is responsible for the fungicidal activity inside the internal tissue of the plant [Buchenauer et al., 1973]. The main key factors for conversing of Thiophanate methyl to biochemical compound inside the plant are the presence of plant enzyme and sunlight [Buchenauer et al., 1973 and Soeda, Kosaka, et al., 1972] (figure 1). Thiophanate methyl transposition to MBC Which is considered a chemically stable metabolite and relatively persistent fungicide, which only metabolized to a limited extent in plants and soil. Thiophanate methyl can be determined in the form of MBC after transformation [Muccio et al., 1995]. Under acidic status the most cost-efficient analytical method that used for control residues of thiophanate-methyl and carbendazim can be shown as the thiophanate-methyl converted to carbendazim by refluxing. UV /VIS spectrophotometer can specify carbendazim [Ono 1973]. This fungicide is applied for protection of several agricultural crops such as grapes, lettuce, citrus fruits, potatoes, cucumber and cereals. Cucumber (*Cucumis sativus*) is the most important cultivated crop belongs to the family of Cucurbitaceae [Heywood, et al., 2007]. It's a creeping vine

vegetables crop. Within cucumber varieties, several cultivars have been created. Cucumber is originally from South Asia, but now it grows in most countries. Many different types of cucumber are traded on the global market. Cucumber is produced in the tropical, subtropical and temperate regions [Heywood et al., 2007 and Judd et al., 2008]. Each 100g of fresh cucumber contains 96g water, 0.1g fat, 2.6-2.4g sugars and minerals. Fungal diseases and insects infections are the main pests that infect cucumber crop which are responsible for yield losses. Utilizing of different pesticides during pest control gives such wonderful results of some time, however the danger of these chemical cannot be ignored due to their residues [Banks, and Soliman 1997, Urani et al., 1995]. The degradation of any complex depends on different factors, including plant species, chemical formulation, climatic conditions, physical phenomenon fundamentally volatilization, application method and chemical degradation, in which sunlight plays a prominent role. Thus, it's important to study the degradation for each crop and fungicides residues under field and protected cultivations appeared that the half-life was 7.2 to 8.3 days in field farming [Abd El-Megeed et al., 2000].



**Figure 1:** Transformation of Thiophanate- methyl carbendazim (MBC)

## MATERIALS AND METHODS

**Chemicals.** HPLC grade is provide from National Center for Pesticide Control (NCPC) /Ministry of Agriculture (Baghdad). Acetonitrile purity (99.9%) from Labscan Ltd. (Ireland); Anhydrous magnesium sulfate  $MgSO_4$  from Sigma Chemical (U.S.A). Sodium chloride NaCl from Avantor (U.S.A). Primary secondary amine (PSA) (40-60  $\mu m$ ), Graphitized Carbon Black (GCB) (200-400 mesh) were from Restek Qsep. (U.S.A) Thiophanate-methyl reference standard was of 99% purity and obtained from Dr. Ehrenstorfer GmbH Germany. Stock solution was prepared in acetonitrile and stored at  $-18^\circ C$ . The standard solutions required for constructing a calibration graph (0.01, 0.25, 0.5, 1, 5, 10 mg/kg) were prepared from stock solution by serial dilution with acetonitrile and were stored at  $4^\circ C$  before use.

**2.2. Field Experiment and Sampling:** A field study was conducted during winter seasons 2018 at Organic Agriculture Research Centre, Plant Protection department, Baghdad: Abu-Ghreb. The cucumber cultivated seeds were obtained by Dr. Enad Daher University of Baghdad College of Agriculture. Soil was tested before the seedling process in the laboratories of the soil department of Agricultural Research as the results were as follows: organic matter; 0.86%, pH 7.6, EC; 6.0ms/cm, texture, sandy Soil, Nitrogen 70.0 ppm, Potassium 592ppm, Phosphorus 24.16ppm.

The experiment was conducted in randomized block design in three replicates (each replicate considered as 25 plant) during 2017-2018. Treatment was carried out by Assltawakul sprayer (Turkey made 5 Lt.) equipped with one nozzle. The thiophanate methyl 70% WP was applied in the recommended dose i.e., 1.5 g L<sup>-1</sup> (T2) then the recommended dose was doubled to 3g L<sup>-1</sup> (T3) respectively and double spray in the recommended dose i.e. 1.5g L<sup>-1</sup> (T4). For the degradation study of thiophanate methyl in cucumber samples were collected periodically by randomly at different time interval zero time (1 h after the application). 1, 3, 5, 7 and 15, 21, 28 days after the application. A 500 mg of cultivated cucumber was collected as sample from each replicate including control line. These samples were transferred by cooler box to the laboratory for storing at ( $-20^\circ C$ ) in freezer.

**2.3 Sampling Preparation:** Samples were performed by randomly collecting from different places of the experimental plots according to the FAO/WHO references. The samples were extracted according to the QuEChERS method described by Muccio et al., [1995]. A 10 g of the homogenized samples from weighted in a 50 ml centrifuge tube and then 10ml of acetonitrile added to the mixture.

The mixture was shaken for 1min using vortex at maximum

Then a 4 g of anhydrous magnesium sulfate ( $MgSO_4$ ) and 1 g of sodium chloride (NaCl) were added to the tube. The mixture was extracted and centrifuged at 3500 rpm for 5 min and  $4^\circ C$ . The matrix was cleaned up by dispersive solid-phase extraction. A 50 mg/ml Primary Secondary Amine (PSA) and 150mg/ml magnesium sulfate ( $MgSO_4$ ) were added to the tube and then the matrix was centrifuged for 10min. at 3500 rpm. The final volume of the matrix was analyzed by HPLC.

**2.4: HPLC Instrument:** A Shimadzu LC-20AD (UFLC) Ultra Fast Liquid Chromatography, High performance liquid chromatography with UV/visible detector was made in Japan for detecting the fungicide residues. The HPLC used reversed phase Kromasil C18 column for Separation the residues from crops. The quality of column was (4.0mm  $\times$  250 mm) with a 5 $\mu m$  particle size (France made, Touzart & Matignon,). To work, the column needs to be thermostat at  $40^\circ C$ . Acetonitrile is Mobile phase, it consists of: water (95:5 v/v). A new pump was used at flow rate of 0.5 mL min<sup>-1</sup> injection volume was 5 $\mu L$ . The wave length was fixed at 230 nm and the run time was 10 min per sample.

**2.5: Standard preparation:** A 0.02 gm of references standard Thiophanate- methyl 99.96% from dr. ehrenstorfer gmbh company was weighted and dissolved of acetonitrile. The mixture was then transferred into a 100 ml volumetric flask and diluted with acetonitrile to the mark to prepare 200 $\mu g/ml$ <sup>-1</sup> stock solutions. The stock solution was used then for preparing of calibration curve and recovery test.

Acetonitrile in order to obtain the standard solutions for spiking and calibration curve. The standard solutions were freshly prepared, filtered through 0.45 $\mu m$  membrane filters, and kept at  $4^\circ C$  in the dark.

**2.5: Recovery Assays:** Un-treated cucumber plant samples were used. These untreated samples were fortified with 0.01, 0.1mg/ml of standard solution. These fortified samples were extracted and clean up and then analyses by HPLC.

**Table 1:** Recoveries and relative standard deviations for Thiophanate-methyl in cucumber at various fortification level.

Fortified level (mg/kg)	Recovery	RSD
0.01	106	2.3
0.05	92	3.2

## RESULTS AND DISCUSSION

Based on a study was conducted by Abd El-Megeed et al., [2000] that the Half Time of Thio-

phanate- methyl was 7.2 to 8.3 days for both cultivated approach greenhouse and open field in addition the average of separation of the fungicide was one application during control pest.

**1.Linearity**, The QuechERS method was used to determine the residue of the Thiophanate - methyl fungicide. It was found that the average recovery rate of all analyzes and external parameters was constant from 92-102% (Figure 3), which indicates the efficiency of this method and allowed identification and comparison of the fungicide in the samples. The values of the relative standard deviation (RSD) were less than 5% (Table 1). In all cases, and generally acceptable for further analysis, the calibration of the quantitative (LOQ) scale was performed by using the external standard calibration curves, which were linear

With a correlation coefficient  $R^2=0.998$  for each analysis. The Limit Of Detection (LOD) level for each analysis in the cucumber samples ranged from 0.001 mg to 1 kg, ensuring that the values were significantly lower than the allowable residue values on the cucumber. Then known concentration of reference standard was added to each sample 0.05, 0.01ml/L. respectively. The fortified samples were analyzed (Fig. 4).

As a result, the Degradation of Thiophanate – methyl was recorded at between 25-27 days.




Figure 3: Calibration curve of Thiophanate-Methyl from 50 to 0.05




Figure 4: Degradation curve of Thiophanate-Methyl

## REFERENCES

- Abd El-Megeed, M.I., Z.H. Zidan, F.A. Afifi and Z.A. Al-Naser, Residues of procymidone and thiophanate-methyl fungicides on tomatoes and cucumber under field and protected cultivations. *Ann. Agric. Sci. Cairo* 4: 1695-1709 (2000).
- Banks D. and M.R. Soliman, Protective effects of antioxidants against benomyl-induced lipid peroxidation and glutathione depletion in rats. *Toxicology* 116: 177–181 (1997).
- Buchenauer H., Erwin D.C. and N.T. Keen, Systemic fungicidal effects of thiophanate methyl on verticillium wilt of cotton and its transformation to methyl-2-benzimidazol carbamate in cotton plants. *Phytopathology*. 63: 1091-1095 (1973).
- Heywood, V.H., Brummitt, R.K., Culham, A. and O Seberg, Flowering Plant Families of the World. Ontario: Firefly Books. In English (2007).
- Judd, W.S., Campbell, C.S., Kellogg, E.A., Stevens, P.F. and M.J. Donoghue, Plant Systematics—A Phylogenetic Approach Third Edition. Sunderland, Massachusetts: Sinauer Associates, Inc. In English (2008)
- Maranghi, F., Macrí, C., Ricciardi, C., Stazi, A.V., Rescia, M. and A. Mantovani, Histological and Histo-morphometric Alterations in Thyroid and Adrenals of CD Rat Pups Exposed in Utero to Methyl Thiophanate. *Reprod. Toxicol.* 17: 617-623 (2003).
- Muccio A.D., Cammoni I., Ventriglia M., Barbini D.A., Mauro M., Pelosi P., Generali T., Ausili A. and S. Girolimetti, Simplified clean-up for the determination of benzimidazolic fungicides by highperformance liquid chromatography with UV detection. *J. Chromatogr A* 697: 145–152 (1995)
- Naqvi S. Tatheer, Asghari Bano, Safia Ahmed, Nisar Ahmed, Irfan Ali and Jahangir Asad, *Pak. J. Biotechnol.* 3 (1-2): 25-32 (2006).
- Ono S., Analytical method for residues of thiophanate-methyl and 2-methyl benzimidazole carbamate in crops by UV (1973).
- Soeda, Y., Kosaka, S. and T. Noguchi, The fate of thiophanate. methyl fungicide and its metabolites on plant leaves and glass plates. *Agr. Biol. Chem.* 36(6): 931-936 (1972).
- Urani C., Chiesara E., Galvani P., Marabini L., Santagostino A. and M. Camatini, Benomyl affects the microtubule cytoskeleton and the glutathione level of mammalian primary cultured hepatocytes. *Toxicol. Lett.* 76: 135–144 (1995).